arXiv:1405.6871v1 [astro-ph.SR] 27 May 2014

The relativistic three dimensional evolution of SN 1987A

L. Zaninetti

Dipartimento di Fisica, via P.Giuria 1,
I-10125 Turin,Italy

E-mail: zaninetti@ph.unito.it

Abstract. The high velocities observed in supernovae require a relativistic treatment for the equation of motion in the presence of gradients in the density of the interstellar medium. The adopted theory is that of the thin layer approximation. The chosen medium is auto-gravitating with respect to an equatorial plane. The differential equation which governs the relativistic conservation of momentum is solved in numerically and by recursion. The asymmetric field of relativistic velocities as well the time dilation are plotted at the age of 1 yr for SN 1987A.

Keywords: supernovae: general supernovae: individual (SN 1987A) ISM : supernova remnants

1. Introduction

The expansion velocities in supernovae (SN) are quite high and, as an example, a time series of eight spectra in SN 2009ig reports that the velocity at the CA II line, decreases from 32000 kms^{-1} to 21500 kms^{-1}, in 12 day, see Fig. 9 in [1]. Another example is given by SN 2009bb in which the velocity of expansion has been evaluated to be ≈ 255000 kms^{-1}, see [2].

We briefly recall that the corrections in special relativity (SR) for stable atomic clocks in satellites of the Global Positioning System (GPS) are applied to satellites which are moving at a velocity of $\approx 3.87\,\mathrm{kms}^{-1}$. The problem of the aspherical SN, such as SN 1987A , is to find an acceptable model which can reproduce the observed complex morphology of the aspherical SN 1987A and this was done in a classical framework by [3]. In this paper we shall discuss a relativistic treatment of the thin layer approximation in the presence of an auto-gravitating medium.

2. Relativistic conservation of momentum

The chosen auto-gravitating profile is

$$n(R,\theta) = n_0 sech^2(\frac{R\sin(\theta)}{2\,h}) \quad , \tag{1}$$

where n_0 is the density in the equatorial plane ($\theta = 0$), R is the radius of the advancing shell, θ is the latitude angle ($\theta = 0$ at the equator and $\theta = \pm 90$ at the two poles) and h is a parameter which characterizes the gradient. The chosen symmetry imposes that the motion is independent of the azimuthal angle in spherical coordinates but depends only on the latitude angle and the time. The classical conservation of momentum in the presence of an auto-gravitating medium was treated in [3] and therefore we will not duplicate the results already obtained. The relativistic conservation of momentum, see [4, 5, 6], is formulated as

$$M(R_0;b)\gamma_0\beta_0 = M(R;b)\gamma\beta \quad , \tag{2}$$

with

$$\gamma_0 = \frac{1}{\sqrt{1-\beta_0^2}} \quad ; \qquad \gamma = \frac{1}{\sqrt{1-\beta^2}} \quad , \tag{3}$$

and

$$\beta_0 = \frac{v_0}{c} \quad ; \qquad \beta = \frac{v}{c} \quad , \tag{4}$$

c being the velocity of light, here $M(R_0;b)$ is a first mass between 0 and R_0 and $M(R;b)$ is a second mass between 0 and R. We know already that $M(R;b) = (I_m(R))^{1/p}$ where the integral $I_m(R)$ has been defined in eq. (15) of [3] and p is a parameter to be found. The fundamental Eq. (2) can be first solved for β^2

$$\beta^2 = \frac{N}{D} \quad , \tag{5}$$

where

$$N = {R_0}^{6\,p^{-1}}{\beta_0}^2 \quad ,$$

$$
\begin{aligned}
D = & -(-1)^{2\,p^{-1}}(-CR0^3S^3 + 8\,C\ln(1+B)Rh^2S \\
& -8\,C\ln(1+A)R0\;h^2S + 4\,CR0^2hS^2 - 4\,CR^2hS^2 \\
& -BR0^3S^3 - AR0^3S^3 + 8\,CP(2,-B)h^3 - 8\,CP(2,-A)h^3 \\
& +8\,B\ln(1+B)Rh^2S - 8\,B\ln(1+A)R0\;h^2S - 4\,BR^2hS^2 \\
& +8\,A\ln(1+B)Rh^2S - 8\,A\ln(1+A)R0\;h^2S + 4\,AR0^2hS^2 \\
& -R0^3S^3 + 8\,BP(2,-B)h^3 - 8\,BP(2,-A)h^3 + 8\,AP(2,-B)h^3 \\
& -8\,AP(2,-A)h^3 + 8\,h^2R\ln(1+B)S - 8\,h^2R0\;\ln(1+A)S \\
& +8\,h^3P(2,-B) - 8\,h^3P(2,-A))^{2\,p^{-1}}(1+B)^{-2\,p^{-1}}S^{-6\,p^{-1}}(1+A)^{-2\,p^{-1}}{\beta_0}^2 \\
& +(-1)^{2\,p^{-1}}(-C{R_0}^3S^3 + 8\,C\ln(1+B)Rh^2S - 8\,C\ln(1+A)R_0\;h^2S \\
& +4\,C{R_0}^2hS^2 - 4\,CR^2hS^2 - B{R_0}^3S^3 - A{R_0}^3S^3 + 8\,CP(2,-B)h^3 \\
& -8\,CP(2,-A)h^3 + 8\,B\ln(1+B)Rh^2S - 8\,B\ln(1+A)R_0\;h^2S - 4\,BR^2hS^2 \\
& +8\,A\ln(1+B)Rh^2S - 8\,A\ln(1+A)R_0\;h^2S + 4\,A{R_0}^2hS^2 - {R_0}^3S^3 \\
& +8\,BP(2,-B)h^3 - 8\,BP(2,-A)h^3 + 8\,AP(2,-B)h^3 - 8\,AP(2,-A)h^3 \\
& +8\,h^2R\ln(1+B)S - 8\,h^2R_0\;\ln(1+A)S + 8\,h^3P(2,-B) \\
& -8\,h^3P(2,-A))^{2\,p^{-1}}(1+B)^{-2\,p^{-1}}S^{-6\,p^{-1}}(1+A)^{-2\,p^{-1}} + {R_0}^{6\,p^{-1}}{\beta_0}^2 \quad ,
\end{aligned}
$$

with

$$
\begin{aligned}
A &= \mathrm{e}^{\frac{R_0\sin(\theta)}{h}} \\
B &= \mathrm{e}^{\frac{R\sin(\theta)}{h}} \\
C &= \mathrm{e}^{\frac{\sin(\theta)(R_0+R)}{h}} \\
S &= \sin(\theta) \quad .
\end{aligned}
$$

and P the polylog operator, which is defined by

$$
polylog(a,z) = \sum_{n=1}^{\infty}\frac{z^n}{n^a} \quad . \tag{6}
$$

The value of β is

$$
\beta = \sqrt{\frac{N}{D}} \quad , \tag{7}
$$

or

$$
\frac{dR}{dt} = c\sqrt{\frac{N}{D}} \quad . \tag{8}
$$

This first order differential equation can be solved with the Runge–Kutta method, see FORTRAN SUBROUTINE RK4 in [7]. Another approach separates the variables

$$
\int_{R_0}^{R}\frac{1}{\sqrt{\frac{N}{D}}}\,dR = c(t-t_0) \quad . \tag{9}
$$

The previous integral does not have an analytical solution and we treat the previous result as a non-linear equation to be solved with the FORTRAN SUBROUTINE ZRIDDR in [7]. The presence of an analytical expression for β as given by Eq. (8) allows setting up the recursive solution

$$
\begin{aligned}
R_{n+1} &= R_n + V_n(R_0, R_n, \beta_0, h)\Delta t \\
V_{n+1} &= V_n(R_0, R_{n+1}, \beta_0, h) \quad ,
\end{aligned} \tag{10}
$$

Table 1. The numerical values of the parameters of the relativistic simulation for SN 1987A

Quantity	Unit	value
R_0	pc	0.011
$\dot{R}_0$	km s^{-1}	30000
p	number	4
h	pc	0.01
t_0	yr	0.00022
t	yr	23

where R_n, V_n, Δt are the temporary radius, the relativistic velocity, and the interval of time, respectively. An interesting application of SR is the time delay: given an interval of time, Δt, in the laboratory frame the interval of time, $\Delta t'$, in a frame that that is moving with velocity v in the x-direction is

$$\Delta t' = \frac{\Delta t}{\sqrt{1 - \frac{v^2}{c^2}}} \quad . \tag{11}$$

We can therefore introduce the following ratio

$$D = \frac{\Delta t}{\Delta t'} \quad , \tag{12}$$

which measures the time dilation, and lies between 0 and 1.

3. Astrophysical application

We numerically solved the non-linear equation, Eq. (9) even if the same results can be obtained by solving the differential equation (8) or implementing the recursive relationship as given by Eq. (10), see Table 1 for the adopted data.

The complicated structure of SN 1987A is due to the great variety of shapes obtained when the point of view of the observer changes. One way to parametrize the point of view of the observer is the introduction of the Euler angles (Φ, Θ, Ψ), as an example, Fig. 1 shows the 3D advancing shell after 23 years. In order to avoid complicated changes of framework for the field of velocity we limit ourselves to the non-rotated image. This choice is already widely used by astronomers in order to reduce the data of η-Carinae , see Fig. 4 in [8]. The progressive increase of the asymmetry is clearly outlined in Fig. 2, in which sections of the expansion are drawn at time steps of 1 yr. The difference in velocity between the polar direction and equatorial direction are oulined in Fig. 3. The relativistic field of velocity in the various points of SN 1987A after 1 yr was shown in Fig. 4. The relativistic time dilation is mapped in Fig. 5 where the velocity of expansion perpendicular to the observer (x-direction) is considered.

4. Conclusions

We have covered the evolution of a SN in an auto-gravitating medium in a relativistic framework. The initial shape is represented by a sphere of radius $R_0 = 0.011$ pc. After 1 yr, the asymmetry between the radius in the equatorial plane and the radius



Figure 1. Continuous three-dimensional surface of SN 1987A after 23 yr: the three Eulerian angles characterizing the point of view are Φ=105°, Θ=55 ° and Ψ=-165 °. Physical parameters as in Table 1.



Figure 2. Sections of SN 1987A in the *X*-*Z* plane at time steps of 1yr. Physical parameters as in Table 1. This is a non-rotated image and the three Euler angles characterizing the orientation are Φ=180°, Θ=90 ° and Ψ=0 °.



Figure 3. Velocity in the equatorial direction (full line) and in the polar direction (dashed line) in the first 50 days.



Figure 4. Map of the relativistic velocity as a function of the latitude for SN 1987A at the age 1 yr.

in the polar direction is well defined and Fig. 4 summarizes both the asymmetrical shape and the anisotropic field of velocity. The time dilation at 1 yr as represented by the parameter D varies between a minimum of 0.9975 and a maximum of 1.

REFERENCES

[1] Marion G H, Vinko J and Wheeler J C 2013 High-velocity Line Forming Regions in the Type Ia Supernova 2009ig *ApJ* **777** 40
[2] Soderberg A M, Chakraborti S and Pignata G 2010 A relativistic type Ibc supernova without a detected γ-ray burst *Nature* **463**, 513 (*Preprint* 0908.2817)
[3] Zaninetti L 2013 Three dimensional evolution of SN 1987a in a self-gravitating disk *International Journal of Astronomy and Astrophysics* **3**, 93
[4] French, AP 1968 *Special Relativity* (New York: CRC)
[5] Zhang Y 1997 *Special Relativity and Its Experimental Foundations* (Singapore: World Scientific)
[6] Guéry-Odelin D and Lahaye T 2010 *Classical Mechanics Illustrated by Modern Physics: 42 Problems with Solutions* (London: Imperial College Press)
[7] Press W H, Teukolsky S A, Vetterling W T and Flannery B P 1992 *Numerical Recipes in*



Figure 5. Map of the relativistic time dilation D, see Eq. (12), for velocity of SN 1987A in the direction perpendicular to the observer at the age 1 yr at time intervals of 1 s in the laboratory frame.

FORTRAN. The Art of Scientific Computing (Cambridge: Cambridge University Press)
[8] Smith N 2006 The Structure of the Homunculus. I. Shape and Latitude Dependence from H2 and Fe II Velocity Maps of eta Carinae *ApJ* **644**, 1151 (*Preprint* `arXiv:astro-ph/0602464`)